

# Supplemental Labeling



Label expiration Date:5/1/2015

EPA Reg. No. 8033-109-279

Active Ingredient:	By Wt.
Acetamiprid .....	5.00 %
Bifenthrin* .....	6.00 %
Other Ingredients: .....	89.00 %
	100.00%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

This product contains 0.44 lb. acetamiprid and 0.53 lb. bifenthrin active ingredients per gallon.

\*Cis isomers 97% minimum, trans isomers 3% maximum.

## DIRECTIONS FOR USE

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

Both this Supplemental labeling and the full Transport Mikron label must be in the possession of the user at the time of application. Read the label affixed to the container for Transport Mikron Insecticide before applying. Use of Transport Mikron Insecticide according to this supplemental labeling is subject to all use precautions and limitations imposed by the labeling affixed to the container for Transport Mikron Insecticide.



FMC Corporation  
Agricultural Products Group  
1735 Market Street  
Philadelphia PA 19103

<b>Food/Feed Handling Establishment Applications</b>	
<b>Controls</b>	Ants (including Red Imported Fire Ants and Carpenter Ants) (but excluding Pharaoh ants and Argentine ants) Bees, Bed Bugs, Bees, Beetles* (including Carpet Beetles) (*Not for use in California), Boxelder Bugs, Centipedes, Cockroaches, Crickets, Earwigs, Firebrats, Fleas, Flies, Gnats, Midges, Millipedes, Moths, (including Cloth Moths), Pillbugs, Scorpions, Silverfish, Sowbugs, Spiders (including Black Widow and Brown Recluse), Springtails, Stink Bugs, Ticks, Wasps
<b>Where to Apply</b>	Applications of this product are permitted in both food/feed and non-food areas of food/feed establishments as a spot or crack and crevice treatment. Food/feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. Including also are areas for receiving, storing, packaging (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, edible oils, syrups) or food. Service areas where food is exposed and the facility is in operation are also considered food areas. Permitted non-food areas of use include, garbage rooms, lavatories, entries and vestibules, offices, locker rooms, machine rooms, garages, mop closets and storage (after canning or bottling). Permitted use sites include: aircraft (do not use in aircraft cabins), apartment buildings, bakeries, bottling facilities, breweries, buses, cafeterias, candy plants, canneries, dairy product processing plants, food manufacturing plants, food processing plants, food service establishments, granaries, grain mills, grocery stores (do not apply directly to the interior of food grain bins) hospitals, hotels, industrial buildings, laboratories, meat/poultry/egg processing plants, mobile/motor homes, nursing homes, offices, pet stores (do not apply directly to the interior of food grain bins or animals), railcars, restaurants, schools, ships, trailers, trucks, vessels, warehouses and wineries.
<b>Application Rate</b>	Apply Transport Mikron Insecticide in a sufficient amount of water (see Dilution Chart) to adequately cover 1,000 square feet. Do not apply more than 1.25 ounces of Transport Mikron Insecticide per 1,000 square feet.
<b>Spot, Crack and Crevice Application</b>	Spot or crack and crevice applications may be made while the facility is in operation; however, cover or remove food from area being treated. Do not apply directly to food.
<b>Mixing Directions</b>	When using spray rigs, fill tank ¼ to 1/3 full with water. Start pump to begin by-pass agitation and place end of treating tool in tank to allow circulation through hose. Add Transport Mikron Insecticide. Add remaining amount of water. Let pump run and allow recirculation through the hose for 2 to 3 minutes. For backpacks and handheld sprayers, fill the tank ¼ full with water. Add Transport Mikron Insecticide. Agitate tank gently before adding remaining water. Close application equipment. For other types of sprayers, Transport Mikron Insecticide may be mixed into full tanks of water. Fill tank with the desired volume of water and add Transport Mikron Insecticide. Close and shake before use to ensure proper mixing. Mix only the amount of dilution needed for application.
<b>Foam Applications</b>	For foaming directions, please refer to FOAM APPLICATIONS FOR CONTROL OF LISTED HOUSEHOLD PESTS in the SPECIFIC PEST CONTROL APPLICATIONS section.

## RESTRICTIONS

Do not apply as a perimeter treatment to areas beyond 10 feet from the foundation of the structure unless using a spot treatment.

Do not use as a space or broadcast spray .

Do not use in and around the exterior perimeter of commercial barns, stables, and paddocks. Do not use in grazing areas, feed lots or other similar areas used for housing, boarding, and/or rearing animals This product may be used around barns and stables on residential property.

Do not apply by air.

Do not apply as a broadcast spray indoors or as a broadcast spray on lawns and turf.

Do not apply in greenhouses or nurseries.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

Do not apply this product through any kind of irrigation system.

Not for use on sod farm turf, golf course turf, or grass grown for seed.

Do not apply to pets, crops, or sources of electricity.

Do not treat electrically active underground services.

Do not treat areas when food is exposed. Cover or remove food from area being treated. Do not allow spray to contact food, foodstuffs, food contacting surfaces, food utensils or water supplies.